

VCS MODULE VMD0001
REDD METHODOLOGICAL MODULE:
ESTIMATION OF CARBON STOCKS IN THE
ABOVE- AND BELOW GROUND BIOMASS IN LIVE
TREE AND NON-TREE POOLS (CP-AB)

Version 1.1

11 October 2013

Sectoral Scope 14

Methodology developed by:



CONTENTS

1	Sources	4
2	Summary Description of the Module	4
3	Definitions and Acronyms	4
4	Applicability Conditions	4
5	Procedures	4
6	Parameters.....	12
6.1	Data and Parameters Available at Validation	12
6.2	Data and Parameters Monitored	18

1 SOURCES

This module is one of numerous modules that comprise the VCS approved methodology VM0007: REDD Methodology Modules.

2 SUMMARY DESCRIPTION OF THE MODULE

This module allows for *ex ante* estimation of carbon stocks in above- and belowground tree and non-tree woody biomass in the baseline case (for both pre- and post-deforestation stocks) and project case and for *ex post* estimation of change in carbon stocks in above- and belowground tree biomass in the project case. Uncertainty of estimates is treated in module X-UNC. Identification of baseline (post-deforestation) land-uses and stocks is treated in modules BL-UP and BL-PL.

3 DEFINITIONS AND ACRONYMS

Definitions

All terms in the following module are used inline with VCS program definitions of the term or, where applicable, the definitions given in VM0007: REDD Methodology Modules.

4 APPLICABILITY CONDITIONS

This module is applicable to all forest types and age classes. Inclusion of the aboveground tree biomass pool as part of the project boundary is mandatory as per the framework module REDD-MF.

Non-tree aboveground biomass must be included as part of the project boundary if the following applicability criteria are met (per framework module REDD-MF):

- Stocks of non-tree aboveground biomass are greater in the baseline than in the project scenario, and
- Non-tree aboveground biomass is determined to be significant (using the T-SIG module).

Belowground (tree and non-tree) biomass are not required for inclusion in the project boundary because omission is conservative.

5 PROCEDURES

Frequency of measurement for baseline above- and belowground biomass stocks

Measurements of initial stocks employed in the baseline must take place within ± 5 years of the project start date, for simplicity referred to here as stocks at $t=0$.

Above- and belowground biomass stock estimates are valid in the baseline (i.e. treated as constant) for 10 years, after which they must be re-estimated from new field measurements. For each stratum, where the re-measured estimate is within the 90% confidence interval of the $t=0$ estimate, the $t=0$ stock estimate takes precedence and is re-employed, and where the re-measured estimate is outside (i.e. greater than or less than) the 90% confidence interval of the $t=0$ estimate, the new stock estimate takes precedence and is used for the subsequent period.

Part 1: Aboveground tree biomass: Estimation of carbon stocks in aboveground tree biomass ($C_{AB_tree,i}$)

The mean carbon stock in aboveground tree biomass per unit area is estimated based on field measurements in sample fixed area plots or sample points using prisms employing representative random or systematic sampling.

Two methods are available for sampling: Fixed Area Plots and Point Sampling with Prisms, both using Allometric Equations method to estimate biomass from measured tree dimensions.

Part 1, Option 1. Fixed Area Plots with Allometric Equation method

Step 1: Determine the tree dimensions such as diameter (*DBH*, at typically 1.3 m [4.3 ft] aboveground level or above buttress where they exist, and total height *H*), of all the trees above some minimum *DBH* in the sample plots. The exact tree dimensions and minimum size tree to be measured in sample plots will be specified by the allometric equation selected in Step 2. Any minimum values employed in inventories are held constant for the duration of the project.

Step 2: Select or develop an appropriate and validated allometric equation for forest type/group of species *j* (e.g. tropical humid forest or tropical dry forest) or for each species or family *j* (group of species) found in the inventory (hereafter referred to as species group). Any equation selected may only be used if applicability has been demonstrated and validated per guidance in the parameters section below.

Step 3: Estimate carbon stock in aboveground biomass for each individual tree of species group *j* in the sample plot located in stratum *i* using the selected or developed allometric equation applied to the tree dimensions resulting from Step 1 and sum the carbon stocks in the sample plot:

$$C_{AB_tree,sp,i} = \sum_j^S \sum_{l=1}^{N_{j,sp,i}} f_j(X, Y...) * CF_j \tag{1}$$

Where:

- $C_{AB_tree,sp,i}$ = Carbon stock in aboveground biomass of trees in plot *sp* in stratum *i*; t C
- CF_j = Carbon fraction of biomass for species group *j*; t C t⁻¹ d.m.
- $f_j(X, Y...)$ = Aboveground biomass of trees based on allometric equation for species group *j* based on measured tree variable(s); t. d.m. tree⁻¹
- i* = 1, 2, 3, ... *M* strata
- j* = 1, 2, 3 ... *S* tree species
- l* = 1, 2, 3, ... $N_{j,sp,i}$ sequence number of individual trees of species group *j* in sample plot *sp* in stratum *i*

Step 4: Calculate the mean carbon stock in aboveground biomass for each stratum, converted to carbon dioxide equivalents:

$$C_{AB_tree,i} = \sum_{sp=1}^{P_i} \frac{C_{AB_tree,sp,i}}{A_{sp,i}} * \frac{44}{12} \tag{2}$$

Where:

- $C_{AB_tree,i}$ = Mean aboveground biomass carbon stock in stratum *i*; t CO₂-e ha⁻¹
- $C_{AB_tree,sp,i,t}$ = Aboveground biomass carbon stock of trees in sample plot *sp* of stratum *i*, t C
- A_{spi} = Area of sample plot *sp* in stratum *i*; ha

sp = 1, 2, 3 ... P_i sample plots in stratum i
 i = 1, 2, 3 ... M strata
 44/12 = Ratio of molecular weight of CO₂ to carbon, t CO₂-e t C⁻¹

Part 1, Option 2: Point Sampling with Allometric Equation method

Step 1: Determine the tree dimensions such as diameter (DBH , at typically 1.3 m [4.3 ft] aboveground level and total height H), of all the trees above some minimum DBH in the sample plots. The exact tree dimensions and minimum size tree to be measured at sample points will be specified by the allometric equation selected in Step 2. Any minimum values employed in inventories are held constant for the duration of the project.

Step 2: Select or develop an appropriate and validated allometric equation for each species group j found in the inventory. Any equation selected may only be used if applicability has been demonstrated and validated per guidance in the parameters section below.

Step 3: Estimate carbon stock in aboveground biomass for each individual tree l of species group j at the sample point located in stratum i using the selected or developed allometric equation applied to the tree dimensions resulting from Step 1, and sum the carbon stocks for the sample point:

$$C_{AB_tree,j,sp,i,t} = \sum_{j=1}^S \sum_{l=1}^{N_{j,sp,i}} \left(\frac{f_j(X,Y...)*CF_j}{(3.1415/10000)*((DBH/100)*D:RAD)^2} \right) \quad (3)$$

Where:

$C_{AB_tree,j,sp,i}$ = Carbon stock in aboveground biomass of trees of species group j at point sp in stratum i ; t C
 CF_j = Carbon fraction of biomass for tree species group j ; t C t⁻¹ d.m.
 $f_j(X,Y...)$ = Allometric equation for species j linking measured tree variable(s) to aboveground biomass of trees; t. d.m. tree⁻¹
 DBH = Diameter at breast height of tree l of species group j at point sp in stratum i at time t , cm
 $D:RAD$ = Ratio of DBH to plot radius, specific to prism Basal Area Factor (BAF) employed in point sampling
 l = 1, 2, 3, ... $N_{j,sp,i}$ sequence number of individual trees of species j at point sp in stratum i
 i = 1, 2, 3, ... M strata
 j = 1, 2, 3 ... S tree species

Step 4: Calculate the mean carbon stock in aboveground biomass for each stratum, converted to carbon dioxide equivalents:

$$C_{AB_tree,i} = \frac{1}{N} * \sum_{sp=1}^{P_i} C_{AB_tree,sp,i} * \frac{44}{12} \quad (4)$$

Where:

$C_{AB_tree,i}$ = Mean aboveground biomass carbon stock in stratum i ; t CO₂-e ha⁻¹

$C_{AB_tree_sp,i}$	= Mean aboveground biomass carbon stock of trees at point sp , in stratum i ; t C
N	= Number of sample points in stratum i ; dimensionless
sp	= 1, 2, 3 ... P_i sample points in stratum i
i	= 1, 2, 3 ... M strata
44/12	= Ratio of molecular weight of CO ₂ to carbon, t CO ₂ -e t C ⁻¹

Part 2: Belowground tree biomass: Estimation of carbon stocks in belowground tree biomass ($C_{BB_tree,i}$)

The mean carbon stock in belowground tree biomass per unit area is estimated based on field measurements in sample fixed area plots or sample points using prisms, employing representative random or systematic sampling. The mean carbon stock in belowground tree biomass per unit area is estimated based on field measurements of aboveground parameters in sample plots. Root to shoot ratios are coupled with the Allometric Equations method Part 1 Options 1 and 2, to calculate belowground from aboveground biomass.

Part 2, Option 1: Fixed area plots with root to shoot ratio

Step 1: Calculate the belowground tree biomass carbon stock for each plot:

The above ground tree biomass carbon stocks in each plot can be estimated by allometric equation method described in Part 1 Option 1.

$$C_{BB_tree,sp,i} = R * C_{AB_tree,sp,i} \quad (5)$$

Where:

$C_{BB_tree,sp,i}$	= Belowground tree biomass carbon stock of trees in plot sp , in stratum i ; t C
$C_{AB_tree,sp,i}$	= Aboveground tree biomass carbon stock of trees in plot sp , in stratum i ; t C
R	= Root to shoot ratio; t root d.m. t ⁻¹ shoot d.m.
i	= 1, 2, 3, ... M strata

Step 2: Calculate the mean belowground tree biomass carbon stock for each stratum, converted to carbon dioxide equivalents:

$$C_{BB_tree,i} = \sum_{sp=1}^{P_i} \frac{C_{BB_tree_sp,i} * \frac{44}{12}}{A_{sp,i}} \quad (6)$$

Where:

$C_{BB_tree,i}$	= Mean belowground tree biomass carbon stock in stratum i ; t CO ₂ -e ha ⁻¹
$C_{BB_tree_sp,i}$	= Mean belowground tree biomass carbon stock of trees in plot sp , in stratum i ; t C
$A_{sp,i}$	= Area of sample plot sp in stratum i ; ha
sp	= 1, 2, 3 ... P_i sample plots in stratum i
i	= 1, 2, 3 ... M strata
44/12	= Ratio of molecular weight of CO ₂ to carbon, t CO ₂ -e t C ⁻¹

Part 2, Option 2. Point Sampling with root to shoot ratio

Step 1: Calculate the belowground tree biomass carbon stock for each point sampled:

The above ground tree biomass carbon stocks in each point sampled can be found using the allometric equation method described in Part 1 Option 2 above.

$$C_{BB_tree,sp,i} = R * C_{AB_tree,sp,i} \quad (7)$$

Where:

$C_{BB_tree,sp,i}$ = Belowground tree biomass carbon stock of trees in point sp , in stratum i ; t C

$C_{AB_tree,sp,i}$ = Aboveground tree biomass carbon stock of trees in point sp , in stratum i ; t C

R = Root to shoot ratio; t root d.m. t⁻¹ shoot d.m.

i = 1, 2, 3, ... M strata

Step 2: Calculate the mean belowground tree biomass carbon stock for each stratum, converted to carbon dioxide equivalents:

$$C_{BB_tree,i} = \frac{1}{N} * \sum_{sp=1}^{P_i} C_{BB_tree_sp,i} * \frac{44}{12} \quad (8)$$

Where:

$C_{BB_tree,i}$ = Mean belowground tree biomass carbon stock in stratum i ; t CO₂-e ha⁻¹

$C_{BB_tree_sp,i}$ = Mean belowground tree biomass carbon stock of trees at point sp , in stratum i ;
t C

N = Number of sample points in stratum i ; dimensionless

sp = 1, 2, 3 ... P_i sample points in stratum i

i = 1, 2, 3 ... M strata

44/12 = Ratio of molecular weight of CO₂ to carbon, t CO₂-e t C⁻¹

Part 3: Aboveground non-tree biomass: Estimation of carbon stocks in aboveground non-tree woody biomass ($C_{AB_nontree,i}$)

The mean carbon stocks in the non-tree aboveground biomass pool per unit area are estimated based on previously published or default data¹ or field measurements. Non-tree woody aboveground biomass pool includes trees smaller than the minimum tree size measured in the tree biomass pool, all shrubs, and all other non-herbaceous live vegetation².

Non-tree vegetation can be sampled using destructive sampling frames and/or, where suitable, in sampling plots in combination with an appropriate allometric equation for shrubs.

Calculate the mean carbon stock in aboveground non-tree biomass for each stratum by adding the mean carbon stock in aboveground biomass calculated using the sampling frame method to the mean carbon stock in aboveground biomass calculated using the allometric equation method.

¹ Where using published or default data these data must be derived from peer-reviewed literature and must be appropriate to the species in the project area or to the geographic region, elevation and precipitation regime in the project area

² Pursuant to AR-WG 21 that the GHG emissions from removal of herbaceous vegetation are insignificant in A/R CDM project activities and therefore these emissions can be neglected in A/R baseline and monitoring methodologies

$$C_{AB_nontree,i} = C_{AB_nontree_sample,i} + C_{AB_nontree_allometric,i} \quad (9)$$

Where:

- $C_{AB_nontree,i}$ = Mean aboveground non-tree biomass carbon stock in stratum i ; t CO₂-e ha⁻¹
 $C_{AB_nontree_sample,i}$ = Mean aboveground non-tree biomass carbon stock in stratum i from sample frame method; t CO₂-e ha⁻¹
 $C_{AB_nontree_allometric,i}$ = Mean aboveground non-tree biomass carbon stock in stratum i from allometric equation method; t CO₂-e ha⁻¹
 i = 1, 2, 3 ... M strata

Part 3, Option 1. Sampling Frame Method:

In a stratum where non-tree vegetation is spatially variable, large frames should be used (e.g. 1-2 m radius circle). Where non-tree vegetation is homogeneous, smaller frames can be used (e.g. 30 cm radius).

Generally, the frame is placed at a randomly or systematically selected GPS point or tree plot. At each location, all vegetation originating from inside the frame is cut at the base and weighed. One representative subsample of the cut material is weighed to obtain its wet mass. The collected subsample is taken to a laboratory, oven dried and weighed to determine the dry mass. The wet to dry ratio of the subsample is then used to estimate the dry mass of the original sample.

To estimate the mean carbon stock per unit area in the aboveground non-tree biomass for each stratum:

$$C_{AB_nontree_sample,i} = \sum_{sfp=1}^{SFP_i} \frac{C_{AB_nontree_sample,sf,i}}{A_{sfp,i}} * CF * \frac{44}{12} \quad (10)$$

Where:

- $C_{AB_nontree_sample,i}$ = Carbon stock in aboveground non-tree vegetation in sampling plot in strata i from sample method; t CO₂-e ha⁻¹
 $C_{AB_nontree_sample,sp,i}$ = Biomass in aboveground non-tree vegetation in sample plot sfp in stratum i from sampling frame method; kg d.m.
 CF_j = Carbon fraction of dominant non-tree vegetation j ; t C t d.m.⁻¹
 $A_{sfp,i}$ = Area of non-tree sampling plot sfp in stratum i ; ha
 sfp = 1, 2, 3 ... SFP_i sample plots in stratum i
 i = 1, 2, 3 ... M strata
 $44/12$ = Ratio of molecular weight of CO₂ to carbon, t CO₂-e t C⁻¹

Part 3, Option 2. Allometric Equation Method:

This method may be used for shrubs, bamboo, or other vegetation types where individuals can be clearly delineated.

Step 1: Select or develop an appropriate allometric equation (if possible species-specific, or if not from a similar species).

Step 2: Estimate carbon stock in aboveground biomass for each individual l in the sample plot r located in stratum i using the selected or developed allometric equation:

$$C_{AB_nontree_allometric,i,r} = \sum_{j=1}^S \sum_{l=1}^{N_{i,r}} f_j(\text{vegetation parameters}) * CF_j \quad (11)$$

Where:

$C_{AB_nontree_allometric,i,r}$ = Carbon stock in aboveground biomass of non-tree sample plot r in stratum i from allometric equation method; t C

CF_j = Carbon fraction of biomass for species j ; t C t⁻¹ d.m.

$f_j(\text{vegetation parameters})$ = Aboveground biomass from allometric equation for species j linking parameters such as stem count, diameter of crown, height, or others ; t. d.m. individual⁻¹

i = 1, 2, 3, ... M strata

r = 1, 2, 3, ... R non-tree allometric method sample plots in stratum i

j = 1, 2, 3 ... S species

l = 1, 2, 3, ... $N_{i,r}$ sequence number of individuals in sample plot r in stratum i

t = 0, 1, 2, 3 ... t^* years elapsed since start of the REDD project activity

Step 3: Calculate the mean carbon stock in aboveground biomass for each stratum, converted to carbon dioxide equivalents:

$$C_{AB_nontree_allometric,i} = \sum_{r=1}^{R_i} \frac{C_{AB_nontree_allometric,r,i}}{Ar_i} * \frac{44}{12} \quad (12)$$

Where:

$C_{AB_nontree_allometric,i}$ = Mean aboveground biomass carbon stock in stratum i from allometric equation method; t CO₂-e ha⁻¹

$C_{AB_nontree_allometric,r,i}$ = Aboveground biomass carbon stock in non-tree vegetation in sample plot r of stratum i from non-tree allometric sample plots, t C

Ar_i = Area of non-tree allometric method sample plot in stratum i ; ha

r = 1, 2, 3 ... R non-tree allometric method sample plots in stratum i

i = 1, 2, 3 ... M strata

44/12 = Ratio of molecular weight of CO₂ to carbon, t CO₂-e t C⁻¹

Part 4: Belowground non-tree biomass: Estimation of belowground carbon stocks in non-tree vegetation ($C_{BB_{nontree,i}}$)

The mean carbon stock in belowground biomass per unit area is estimated based on field measurements of aboveground parameters in sample plots. Root to shoot ratios are coupled with the aboveground biomass estimate to calculate belowground from aboveground biomass.

Step 1: Select an appropriate root to shoot ratio for non-tree biomass.

Step 2: Use the appropriate root to shoot ratio to estimate the belowground biomass from aboveground biomass carbon stock in non-tree vegetation in sample plot sp of stratum i , t C:

$$C_{BB_{nontree,i,sp}} = C_{AB_{nontree,i,sp}} * R \quad (13)$$

Where:

$C_{AB_{nontree,i,sp}}$ = Aboveground biomass carbon stock in non-tree vegetation in sample plot sp of stratum i , t C

$C_{BB_{nontree,i,sp,t}}$ = Belowground biomass carbon stock in non-tree vegetation in sample plot sp of stratum i , t C

R = Root to shoot ratio; t root d.m. t⁻¹ shoot d.m.

i = 1, 2, 3 ... M strata

sp = 1, 2, 3 ... P_i sample plots in stratum i

Step 3: Calculate the mean carbon stock in belowground biomass for each stratum, converted to carbon dioxide equivalents:

$$C_{BB_{nontree,i}} = \sum_{sp=1}^{P_i} \frac{C_{BB_{nontree,sp,i}}}{A_{sp,i}} * \frac{44}{12} \quad (14)$$

Where:

$C_{BB_{nontree,i}}$ = Mean belowground biomass carbon stock in stratum i ; t CO₂-e ha⁻¹

$C_{BB_{nontree,i,sp}}$ = Belowground biomass carbon stock in non-tree vegetation in sample plot sp of stratum i ; t C

A_{spi} = Area of sample plot sp in stratum i ; ha

sp = 1, 2, 3 ... P_i sample plots in stratum i

i = 1, 2, 3 ... M strata

44/12 = Ratio of molecular weight of CO₂ to carbon, t CO₂-e t C⁻¹

6 PARAMETERS

6.1 Data and Parameters Available at Validation

Data / Parameter:	<i>CF</i>
Data unit	t C t d.m. ⁻¹
Description	Carbon fraction of dry matter in t C t ⁻¹ d.m.
Equations	1,3,10,11
Source of data	Values from the literature (e.g. IPCC 2006 INV GLs AFOLU Chapter 4 Table 4.3) shall be used if available, otherwise default value of 0.47 t C t ⁻¹ d.m. can be used
Value applied	
Justification of choice of data or description of measurement methods and procedures applied	
Purpose of Data	
Comments	Where new species are encountered in the course of monitoring, new carbon fraction values must be sourced from the literature or otherwise use the default value.

Data / Parameter	<i>D:RAD</i>																		
Data unit	Dimensionless																		
Description	Ratio of DBH to plot radius, specific to prism Basal Area Factor (BAF) employed in point sampling																		
Equations	3																		
Source of data	<table border="1"> <thead> <tr> <th>BAF (m²/ha)</th> <th>D:RAD</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>35.4</td> </tr> <tr> <td>3</td> <td>28.9</td> </tr> <tr> <td>4</td> <td>25.0</td> </tr> <tr> <td>5</td> <td>22.4</td> </tr> <tr> <td>6</td> <td>20.4</td> </tr> <tr> <td>7</td> <td>18.9</td> </tr> <tr> <td>8</td> <td>17.7</td> </tr> <tr> <td>9</td> <td>16.7</td> </tr> </tbody> </table>	BAF (m ² /ha)	D:RAD	2	35.4	3	28.9	4	25.0	5	22.4	6	20.4	7	18.9	8	17.7	9	16.7
BAF (m ² /ha)	D:RAD																		
2	35.4																		
3	28.9																		
4	25.0																		
5	22.4																		
6	20.4																		
7	18.9																		
8	17.7																		
9	16.7																		
Value applied:																			

Justification of choice of data or description of measurement methods and procedures applied	
Purpose of Data	
Comments	

Data / Parameter	$f_j(X, Y)$
Data unit	t d.m. tree ⁻¹
Description	Allometric equation for species <i>j</i> linking measured tree variable(s) to aboveground biomass of living trees, expressed as t d.m. tree ⁻¹
Equations	Error! Reference source not found. , 3
Source of data	<p>Equations must have been derived using a wide range of measured variables (e.g. DBH, Height, etc.) based on datasets that comprise at least 30 trees. Equations must be based on statistically significant regressions and must have an r^2 that is ≥ 0.8.</p> <p>The source of equation(s) shall be chosen with priority from higher to lower preference, as available, as follows:</p> <ul style="list-style-type: none"> (a) National species-, genus-, family-specific; (b) Species-, genus-, family-specific from neighboring countries with similar conditions (i.e. broad continental regions); (c) National forest-type specific; (d) Forest-type specific from neighboring countries with similar conditions (i.e. broad continental regions); (e) Pan-tropical forest type-specific <p>such as those provided Tables 4.A.1 to 4.A.3 of the GPG-LULUCF (IPCC 2003) or in</p> <p>Pearson, T., Walker, S. and Brown, S. 2005. Sourcebook for Land Use, Land-Use Change and Forestry Projects. Winrock International and the World Bank Biocarbon Fund. 57pp. Available at: http://www.winrock.org/Ecosystems/files/Winrock-BioCarbon_Fund_Sourcebook-compressed.pdf</p> <p>or in</p> <p>Chave, J., C. Andalo, S. Brown, M. A. Cairns, J. Q. Chambers, D. Eamus, H. Folster, F. Fromard, N. Higuchi, T. Kira, J.-P. Lescure, B. W. Nelson, H. Ogawa, H. Puig, B. Riera, T. Yamakura. 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. <i>Oecologia</i> 145: 87-99.</p> <p>Species-, genus- and family-specific allometric equations may not always be available, and may be difficult to apply with certainty in the typically species rich forests of the humid tropics, hence it is acceptable practice to use equations developed for regional or</p>

	<p>panropical forest types, provided that their accuracy has been validated with direct site-specific data (per guidance below). If a forest-type specific equation is used, it should not be used in combination with species-specific equation(s) (i.e. it must be used for all tree species³).</p>
Value applied:	
Justification of choice of data or description of measurement methods and procedures applied	
Purpose of Data	
Comments	<p>It is necessary to validate the applicability of equations used. Source data from which equation was derived should be reviewed and confirmed to be representative of the forest type/species and conditions in the project and covering the range of potential independent variable values.</p> <p>Allometric equations can be validated either by:</p> <p>1. Limited Measurements</p> <ul style="list-style-type: none"> • Select at least 30 trees (if validating forest type-specific equation, selection should be representative of the species composition in the project area, i.e. species representation in roughly in proportion to relative basal area). Minimum diameter of measured trees shall be 20cm and maximum diameter shall reflect the largest trees present or potentially present in the future in the project area (and/or leakage belt) • Measure DBH, and height to a 10 cm diameter top or to the first branch. • Calculate stem volume from measurements and multiplying by species-specific density to gain biomass of bole. • Apply a biomass expansion factor to estimate total aboveground biomass from stem biomass. For broadleaf tropical trees this factor shall be: <ul style="list-style-type: none"> ○ 1.38 for trees 20-40cm ○ 1.33 for trees 40-80cm ○ 1.25 for trees $\geq 80\text{cm}^4$ • Plot all the estimated biomass of all the measured trees along with the curve of biomass against diameter as predicted by the allometric equation. If the estimated biomass of the measured trees are distributed both above and below the curve (as predicted by the allometric equation) the equation may be used. The equation may also be used if the measured individuals have a biomass consistently higher than predicted by the equation. If plotting the biomass of the measured trees indicates a systematic bias to overestimation of biomass (>75% of the

³ Note that forest type specific and panropical equations will typically not include palm species or hollow-stem species (e.g. Cecropia) and so specific equations for these growth forms will be needed

⁴ Biomass expansion factors conservatively selected from destructively sampled trees in Bolivia and Belize, reference to IPCC GPG LULUCF BEF values and expert opinion.

	<p>trees above the predicted curve) then destructive sampling must be undertaken, or another equation selected.</p> <p>or</p> <p>2. Destructive Sampling</p> <ul style="list-style-type: none"> • Select at least five trees (if validating forest type-specific equation, selection should be representative of the species composition in the project area, i.e. species representation in roughly in proportion to relative basal area) at the upper end of the range of independent variable values existing in the project area • Measure DBH and commercial height and calculate volume using the same procedures/equations used to generate commercial volumes to which BCEFs will be applied • Fell and weigh the aboveground biomass to determine the total (wet) mass of the stem, branch, twig, leaves, etc. Extract and immediately weigh subsamples from each of the wet stem and branch components, followed by oven drying at 70 degrees C to determine dry biomass; • Determine the total dry weight of each tree from the wet weights and the averaged ratios of wet and dry weights of the stem and branch components. • Plot the biomass of all the harvested trees along with the curve of biomass against diameter as predicted by the allometric equation. If the biomass of the harvested trees are distributed both above and below the curve (as predicted by the allometric equation) the equation may be used. The equation may also be used if the harvested individuals have a biomass consistently higher than predicted by the equation. If plotting the biomass of the harvested trees indicates a systematic bias to overestimation of biomass (>75% of the trees below the predicted curve) then additional destructive sampling must be undertaken, or another equation selected. <p>Details of destructive sampling measurements are given in:</p> <p>Brown, S. 1997. Estimating biomass and biomass change of tropical forests: a primer. FAO Forestry Paper 134, Rome, Italy.</p> <p>Available at http://www.fao.org/docrep/W4095E/W4095E00.htm</p> <p>If using species-specific equations, and new species are encountered in the course of monitoring, new allometric equations must be sourced from the literature and validated, if necessary, as per requirements and procedures above.</p>
--	--

Data / Parameter	$f_i(\text{vegetation parameters})$
Data unit	t. d.m. individual ⁻¹

Description	Allometric equation for non-tree species / linking parameters such as stem count, diameter of crown, height, or others to aboveground biomass of an individual
Equations	Error! Reference source not found.1
Source of data	Whenever available, use allometric equations that are species-specific or group of species-specific, provided the equations have been derived using a wide range of diameters and heights, based on datasets that comprise at least 30 individuals. Project participants may create project location specific equation where appropriate.
Value applied:	
Justification of choice of data or description of measurement methods and procedures applied	<p>It is necessary to verify the applicability of existing equations used. Allometric equations can be verified either by:</p> <ol style="list-style-type: none"> 1. Review of source data from which equation was derived and confirmation that the source data is representative of the species and conditions in the project and covers the range of potential sizes. <p>Or</p> <ol style="list-style-type: none"> 2. Destructive Sampling <ul style="list-style-type: none"> • Selecting at least five individuals covering the range of sizes existing, and felling and weighing the aboveground biomass to determine the total (wet) mass of the stem and branch components; • Extracting and immediately weighing subsamples from each of the wet stem and branch components, followed by oven drying at 70°C to determine dry biomass; • Determining the total dry weight of each individual from the wet weights and the averaged ratios of wet and dry weights of the stem and branch components. <p>If the biomass of the harvested individual is within $\pm 10\%$ of the mean values predicted by the selected allometric equation, and is not biased, then mean values from the equation may be used. Otherwise, the equation must be re-parameterized to conform to the validation data before using, or another equation selected.</p> <p>To create a new allometric equation:</p> <p>Follow guidance in: Pearson, T., Walker, S. and Brown, S. 2005. Sourcebook for Land Use, Land-Use Change and Forestry Projects. Winrock International and the World Bank Biocarbon Fund. 57pp. Available at: http://www.winrock.org/Ecosystems/files/Winrock-BioCarbon_Fund_Sourcebook-compressed.pdf</p>
Purpose of Data	
Comments	

Data / Parameter	<i>R</i>																																			
Data unit	t root d.m. t ⁻¹ shoot d.m.																																			
Description	Root to shoot ratio appropriate to species or forest type / biome; note that as defined here, root to shoot ratio is applied as belowground biomass per unit area:aboveground biomass per unit area (not on a per stem basis)																																			
Equations	5,7,13																																			
Source of data	<p>The source of data shall be chosen with priority from higher to lower preference as follows:</p> <p>(a) Detailed data collected using common practices for root sampling in the area;</p> <p>(b) Globally forest type-specific or eco-region-specific (e.g. IPCC GPG-LULUCF).</p> <p><i>Root to shoot ratios for tropical and subtropical forests modified from</i></p> <p><i>Table 4.4. in IPCC GL AFOLU</i></p> <table border="1"> <thead> <tr> <th>Domain</th> <th>Ecological Zone</th> <th>Aboveground biomass</th> <th>Root-to-shoot ratio</th> <th>Range</th> </tr> </thead> <tbody> <tr> <td rowspan="4">Tropical</td> <td rowspan="2">Tropical rainforest</td> <td><125 t.ha⁻¹</td> <td>0.20</td> <td>0.09-0.25</td> </tr> <tr> <td>>125 t.ha⁻¹</td> <td>0.24</td> <td>0.22-0.33</td> </tr> <tr> <td rowspan="2">Tropical dry forest</td> <td><20 t.ha⁻¹</td> <td>0.56</td> <td>0.28-0.68</td> </tr> <tr> <td>>20 t.ha⁻¹</td> <td>0.28</td> <td>0.27-0.28</td> </tr> <tr> <td rowspan="4">Subtropical</td> <td rowspan="2">Subtropical humid forest</td> <td><125 t.ha⁻¹</td> <td>0.20</td> <td>0.09-0.25</td> </tr> <tr> <td>>125 t.ha⁻¹</td> <td>0.24</td> <td>0.22-0.33</td> </tr> <tr> <td rowspan="2">Subtropical dry forest</td> <td><20 t.ha⁻¹</td> <td>0.56</td> <td>0.28-0.68</td> </tr> <tr> <td>>20 t.ha⁻¹</td> <td>0.28</td> <td>0.27-0.28</td> </tr> </tbody> </table>	Domain	Ecological Zone	Aboveground biomass	Root-to-shoot ratio	Range	Tropical	Tropical rainforest	<125 t.ha ⁻¹	0.20	0.09-0.25	>125 t.ha ⁻¹	0.24	0.22-0.33	Tropical dry forest	<20 t.ha ⁻¹	0.56	0.28-0.68	>20 t.ha ⁻¹	0.28	0.27-0.28	Subtropical	Subtropical humid forest	<125 t.ha ⁻¹	0.20	0.09-0.25	>125 t.ha ⁻¹	0.24	0.22-0.33	Subtropical dry forest	<20 t.ha ⁻¹	0.56	0.28-0.68	>20 t.ha ⁻¹	0.28	0.27-0.28
Domain	Ecological Zone	Aboveground biomass	Root-to-shoot ratio	Range																																
Tropical	Tropical rainforest	<125 t.ha ⁻¹	0.20	0.09-0.25																																
		>125 t.ha ⁻¹	0.24	0.22-0.33																																
	Tropical dry forest	<20 t.ha ⁻¹	0.56	0.28-0.68																																
		>20 t.ha ⁻¹	0.28	0.27-0.28																																
Subtropical	Subtropical humid forest	<125 t.ha ⁻¹	0.20	0.09-0.25																																
		>125 t.ha ⁻¹	0.24	0.22-0.33																																
	Subtropical dry forest	<20 t.ha ⁻¹	0.56	0.28-0.68																																
		>20 t.ha ⁻¹	0.28	0.27-0.28																																
Value applied:																																				
Justification of choice of data or description of measurement methods and procedures applied																																				
Purpose of Data																																				

Comments	<p>Guidelines for Conservative Choice of Default Values:</p> <ol style="list-style-type: none"> 1. If in the sources of data mentioned above, default data are available for conditions that are similar to the project (similar forest or vegetation type; same climate zone), then mean values of default data may be used and considered conservative. 2. Global values may be selected from Table 4.4 (modified as given above) of the AFOLU Guidelines (IPCC 2006), by choosing a climatic zone and forest type that most closely matches the project circumstances.
----------	---

6.2 Data and Parameters Monitored

Data / Parameter	A_{sp}
Data unit	ha
Description	Area of sample plots in ha
Equations	2,6,14
Source of data	Recording and archiving of number and size of sample plots
Description of measurement methods and procedures to be applied	
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	
Purpose of data	
Calculation method	
Comments	Where carbon stock estimation occurs only for determination of the baseline this parameter shall be known <i>ex-ante</i> . Where part of project monitoring, <i>ex-ante</i> the number and area of sample plots shall be estimated based on projected sample effort relative to projections of growth and emissions.

Data / Parameter	N
Data unit	Dimensionless
Description	Number of sample points

Equations	4,8
Source of data	Recording and archiving of number of sample points
Description of measurement methods and procedures to be applied	
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	
Purpose of data	
Comments	Where carbon stock estimation occurs only for determination of the baseline this parameter shall be known <i>ex-ante</i> . Where part of project monitoring, <i>ex-ante</i> the number of sample plots shall be estimated based on projected sample effort relative to projections of growth and emissions.

Data / Parameter	<i>DBH</i>
Data unit	cm
Description	Diameter at breast height of a tree in cm
Equations	1,3
Source of data	Field measurements in sample plots
Description of measurement methods and procedures to be applied	Typically measured 1.3m aboveground. Measure all trees above some minimum <i>DBH</i> in the sample plots. The minimum <i>DBH</i> varies depending on tree species and climate; for instance, the minimum <i>DBH</i> may be as small as 2.5 cm or as high as 20 cm, but for himud tropical forests 10 cm is commonly used. Minimum <i>DBH</i> employed in inventories is held constant for the duration of the project.
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	Standard quality control / quality assurance (QA/QC) procedures for forest inventory including field data collection and data management shall be applied. Use or adaptation of QA/QCs already applied in national forest monitoring, or available from

	published handbooks, or form the <i>IPCC GPG LULUCF 2003</i> , is recommended.
Purpose of data	
Calculation method	
Comments	Where carbon stock estimation occurs only for determination of the baseline this parameter shall be known <i>ex-ante</i> . Where part of project monitoring, <i>ex-ante</i> DBH shall be estimated based on projections of growth.

Data / Parameter	A_{sf}
Data unit	m ²
Description	Area of one sampling frame
Equations	10
Source of data	Recording and archiving size of sampling frame plot
Description of measurement methods and procedures to be applied	
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	
Purpose of data	
Calculation method	
Comments	Shall be known <i>ex-ante</i> .

Data / Parameter	A_r
Data unit	Hectares
Description	Total area of all non-tree allometric method sample plots in stratum <i>i</i>
Equations	12
Source of data	Recording and archiving size of non-tree allometric method sample plot
Description of measurement methods	

and procedures to be applied	
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	
Purpose of data	
Calculation method	
Comments	Where carbon stock estimation occurs only for determination of the baseline this parameter shall be known <i>ex-ante</i> . Where part of project monitoring, <i>ex-ante</i> the number and area of sample plots shall be estimated based on projected sample effort relative to projections of growth and emissions.

Data / Parameter	<i>H</i>
Data unit	m
Description	Total height of tree
Equations	1,3
Source of data	Field measurements in sample plots
Description of measurement methods and procedures to be applied	
Frequency of monitoring/recording	Monitoring must occur at least every ten years for baseline renewal. Where carbon stock enhancement is included monitoring shall occur at least every five years
QA/QC procedures to be applied	
Purpose of data	
Calculation method	
Comments	Where carbon stock estimation occurs only for determination of the baseline this parameter shall be known <i>ex-ante</i> . Where part of project monitoring, <i>ex-ante</i> height shall be estimated based on projections of growth.

DOCUMENT HISTORY

Version	Date	Comment
v1.0	3 Dec 2010	Initial version released
v 1.1	10 Oct 2013	Typographic error in equation 10 has been corrected.